

RB Health (US) LLC

-PE Product Labels –

Tamper evident: Do not use if printed seal over cap is torn or missing.

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

Mucinex[®]
SINUS-MAX[®]

3/4 FL OZ (22 mL) + 16 LIQUID GELS (Liquid Filled Capsules)

VALUE PACK! 2 formats in 1 (Nasal Spray & Liquid Gels)

Mucinex[®]
SINUS-MAX[®]

SEVERE NASAL CONGESTION RELIEF

CLEAR & COOL

NASAL SPRAY
Oxymetazoline HCl 0.05%
Nasal Decongestant

✓ Fast acting
✓ Lasts 12 Hours
✓ Cooling Menthol flavor

FOR AGES 6+

3/4 FL OZ (22 mL) + 16 LIQUID GELS (Liquid Filled Capsules)

MAXIMUM STRENGTH

NDC 63824-509-02

Mucinex[®]
SINUS-MAX[®]

PRESSURE, PAIN & COUGH

Acetaminophen – Pain Reliever
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant • Phenylephrine HCl – Nasal Decongestant

✓ Relieves Sinus Pressure & Congestion
✓ Relieves Headache
✓ Controls Cough
✓ Thins & Loosens Mucus

Actual Size

FOR AGES 12+

Maximum Strength per 4-hour dose
Take only as directed.
Please refer to dosing directions in the individual Drug Facts labels. Do not take more than one product at any given time. Ask a doctor or pharmacist before use if you are unsure or have questions about taking these products.

KEEP CARTONS FOR FULL INFORMATION.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2020 RB Health
Made in China

3 63824 12515 2

PRINT SAFE AREA

KO

Mucinex Sinus-Max Pressure, Pain & Cough**Drug Facts****Active ingredients
(in each liquid gel)****Purposes**

Acetaminophen 325 mg.....Pain reliever
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, shellac, sorbitol sorbitan solution, titanium dioxide, water

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

Mucinex Sinus-Max Severe Nasal Congestion Relief**Drug Facts****Active ingredient****Purpose**

Oxymetazoline hydrochloride 0.05%.....Nasal decongestant

Uses

- temporarily relieves nasal congestion due to:
 - a cold
 - hay fever or other upper respiratory allergies
- promotes nasal and sinus drainage
- temporarily relieves sinus congestion and pressure
- helps clear nasal passages; shrinks swollen membranes

Warnings

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland

When using this product

- do not exceed recommended dosage
- do not use this product for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- this product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge
- the use of this container by more than one person may spread infection

Stop use and ask a doctor if symptoms persist

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril, not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: consult a doctor. Shake well before use.
- To open, hold by the white grips then squeeze and turn cap. Before using for the first time, remove the protective cap from the tip and prime metered pump by depressing firmly several times. To spray, hold bottle with thumb at the base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress pump all the way down with a firm even stroke and sniff deeply. Wipe nozzle clean after use. To close, turn cap until it "clicks".

Mucinex Sinus-Max Severe Nasal Congestion Relief (cont'd)**Drug Facts (continued)**

DO NOT DISCARD CAP

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

benzalkonium chloride, camphor, edetate disodium, eucalyptol, glycine, menthol, polyethylene glycol, polysorbate 80, propylene glycol, purified water, sodium chloride, sodium hydroxide

Questions?

1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this number.

To open: hold by the white grips then squeeze & turn cap.



KEEP CARTONS FOR FULL INFORMATION.



HEALTH • HYGIENE • HOME

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
© 2020 RB Health
Made in India

Please visit our website
www.mucinex.com
Patents: www.rb.com/patents

3134411

061220

PRINT SAFE AREA

KO

Drug Facts (continued)

Other information

- each 20 mL contains sodium 12 mg (Fast-Max Cold, Flu & Sore Throat only) and sodium 16 mg (Nightshift Cold & Flu only)
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients (Mucinex Fast-Max Cold, Flu & Sore Throat)

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propyl galate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum
*may contain this ingredient

Inactive ingredients (Mucinex Nightshift Cold & Flu)

ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Do not take Mucinex Fast-Max Cold, Flu & Sore Throat and Mucinex Nightshift Cold & Flu at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid. TAKE ONLY AS DIRECTED.

DAY & NIGHT COLD & FLU RELIEF

MAXIMUM STRENGTH

NDC 63824-119-66

Mucinex
FAST-MAX

COLD, FLU & SORE THROAT

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectoant
Phenylephrine HCl – Nasal Decongestant

- ✓ Controls Cough, Thins & Loosens Mucus
- ✓ Nasal & Chest Congestion
- ✓ Sinus Pressure & Congestion
- ✓ Body Pain, Headache, Fever & Sore Throat

Mucinex
NIGHTSHIFT

COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Triprolidine HCl – Antihistamine



- ✓ COUGH ✓ FEVER ✓ SORE THROAT
- ✓ RUNNY NOSE ✓ SNEEZING

FOR AGES 12+ **TWO – 6 FL OZ (180 mL) bottles TOTAL – 12 FL OZ (360 mL)**

FOR AGES 12+

No coating

Dist. by: RB Health (US), Parsippany, NJ 07054-0224 Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Drug Facts (continued)

Active ingredients (in each 20 mL) Purposes

Mucinex Nightshift Cold & Flu
Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Triprolidine HCl 2.5 mg.....Antihistamine

Drug Facts

Active ingredients (in each 20 mL) Purposes

Mucinex Fast-Max Cold, Flu & Sore Throat
Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectoant
Phenylephrine HCl 10 mg.....Nasal decongestant

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers (Nightshift Cold & Flu only)
- When using this product:
 - do not use more than directed
 - excitability may occur, especially in children (Nightshift Cold & Flu only)
 - marked drowsiness may occur (Nightshift Cold & Flu only)
 - alcohol, sedatives, and tranquilizers may increase drowsiness (Nightshift Cold & Flu only)
 - avoid alcoholic drinks (Nightshift Cold & Flu only)
 - use caution when driving a motor vehicle or operating machinery (Nightshift Cold & Flu only)
- Stop use and ask a doctor if:
 - nervousness, dizziness, or sleeplessness occur (Fast-Max Cold, Flu & Sore Throat only)
 - pain, nasal congestion, or cough gets worse or lasts more than 7 days (Fast-Max Cold, Flu & Sore Throat only)
 - pain, nasal congestion, or cough gets worse or lasts more than 7 days (Nightshift Cold & Flu only)
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back, or occurs with fever, rash, or headache that lasts. The use could be signs of a serious condition.

Drug Facts (continued)

Uses **MUCINEX FAST-MAX COLD, FLU & SORE THROAT**
Temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion
- sore throat
- sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

MUCINEX NIGHTSHIFT COLD & FLU

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - itchy of the nose or throat
 - itchy, watery eyes due to hay fever
 - temporarily reduces fever
 - controls cough to help you get to sleep

Warnings

- Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 - more than 4000 mg in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product
- Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin redness
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease (Fast-Max Cold, Flu & Sore Throat only)
- diabetes (Fast-Max Cold, Flu & Sore Throat only)
- high blood pressure (Fast-Max Cold, Flu & Sore Throat only)
- thyroid disease (Fast-Max Cold, Flu & Sore Throat only)
- glaucoma (Nightshift Cold & Flu only)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (Nightshift Cold & Flu only)





rb
HEALTH • HYGIENE • HOME

Questions?
1-866-MUCINEX

PARENTS:



<p>Drug Facts</p> <p>Active ingredients (in each caplet)</p> <p>Mucinex FAST-MAX DAY TIME SEVERE CONGESTION & COUGH</p> <p>Dextromethorphan HBr 10 mg.....Cough suppressant Guaifenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p>Active ingredients (in each caplet)</p> <p>Mucinex FAST-MAX NIGHT TIME COLD & FLU</p> <p>Acetaminophen 325 mg.....Pain reliever/</p>	<p>Drug Facts (continued)</p> <p>temporarily relieves these common cold and flu symptoms (NIGHT TIME only):</p> <ul style="list-style-type: none"> cough minor aches and pains headache nasal congestion sore throat runny nose sneezing itching of the nose or throat itchy, watery eyes due to hay fever temporarily reduces fever (NIGHT TIME only) controls cough to help you get to sleep <p>Warnings</p> <p>Liver warning (NIGHT TIME only): This product contains acetaminophen. Severe liver damage may occur if you take:</p>	<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. (NIGHT TIME only) with any other product containing diphenhydramine, even one used on the skin (NIGHT TIME only) if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin (NIGHT TIME only) taking sedatives or tranquilizers (NIGHT TIME only) <p>When using this product</p> <ul style="list-style-type: none"> do not use more than directed excitability may occur, especially in children (NIGHT TIME only)
---	--	--	---

<p>Acetaminophen 325 mg Pain reliever/fever reducer Diphenhydramine HCl 12.5 mg Antihistamine/cough suppressant Phenylephrine HCl 5 mg Nasal decongestant</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves (DAY TIME only): <ul style="list-style-type: none"> cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants the intensity of coughing nasal congestion due to a cold helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (DAY TIME only) 	<ul style="list-style-type: none"> more than 12 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks daily while using this product <p>Allergy alert (NIGHT TIME only): Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin reddening blisters rash If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning (NIGHT TIME only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	<p>2 weeks after stopping the MAUI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease (NIGHT TIME only) heart disease diabetes high blood pressure thyroid disease trouble urinating due to an enlarged prostate gland glaucoma (NIGHT TIME only) a breathing problem such as emphysema or chronic bronchitis (NIGHT TIME only) 	<p>marked drowsiness may occur (NIGHT TIME only)</p> <ul style="list-style-type: none"> alcohol, sedatives, and tranquilizers may increase drowsiness (NIGHT TIME only) avoid alcoholic drinks (NIGHT TIME only) be careful when driving a motor vehicle or operating machinery (NIGHT TIME only) <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> nervousness, dizziness, or sleeplessness occur symptoms do not get better within 7 days or occur with fever (DAY TIME only)
<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> pain, nasal congestion, or cough gets worse or lasts more than 7 days (NIGHT TIME only) fever gets worse or lasts more than 3 days (NIGHT TIME only) redness or swelling is present (NIGHT TIME only) new symptoms occur (NIGHT TIME only) cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Overdose warning (NIGHT TIME only): Taking more than the recommended dose (overdose) may cause liver damage. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed (see Overdose warning) do not take more than 12 caplets in any 24-hour period 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> adults and children 12 years of age and over: take 2 caplets every 4 hours children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> store at 20-25°C (68-77°F) <p>Inactive ingredients (Mucinex FAST-MAX DAY TIME SEVERE CONGESTION & COUGH) croscarmellose sodium, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, methacrylic acid-ethyl acrylate copolymer, mica, microcrystalline cellulose, polyethylene glycol 3350, polysorbate 80, polyvinyl alcohol, povidone K29/32, sodium bicarbonate, talc, titanium dioxide</p> <p>Inactive ingredients (Mucinex FAST-MAX NIGHT TIME COLD & FLU) croscarmellose sodium, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, ferric oxide yellow, methacrylic acid-ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose,</p>	<p>Drug Facts (continued)</p> <p>polyethylene glycol, polyvinyl alcohol, povidone, sodium bicarbonate, talc, titanium dioxide</p> <p>Questions? 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.</p>	<p>PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION</p> <p>Do not take the Mucinex FAST-MAX DAY TIME SEVERE CONGESTION & COUGH and Mucinex FAST-MAX NIGHT TIME COLD & FLU caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. Do not take more than a total of 12 caplets in any 24-hour period.</p> <p>Take only as directed.</p> <p> HEALTH • HYGIENE • HOME</p> <p>3091262 071519</p> <p></p>



8326924

020618

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

3 "63824"55130" 2

Tamper evident: do not use if outer package
is opened or if blister is torn or broken.www.mucinex.com

8326925

Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224 Made in England ©2018 RB 020218

Drug Facts (continued)

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison

Drug Facts (continued)**Other information**

- store between 20-25°C (68-77°F)

Inactive ingredients
(Mucinex FAST-MAX
DAY TIME COLD & FLU)

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients
(Mucinex FAST-MAX
PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION

Do not take the Mucinex Fast-Max DAY TIME COLD & FLU and Mucinex Fast-Max NIGHT TIME COLD & FLU caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. Do not take more than a total of 12 caplets in any 24-hour period. Take only as directed. 020618 8326926

Drug Facts

Active ingredients (in each caplet)	Purposes
Mucinex FAST-MAX DAY TIME COLD & FLU	
Acetaminophen 325 mg	

Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

NIGHT TIME COLD 2887)
 croscarmellose sodium, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, ferric oxide yellow, methacrylic acid – ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium bicarbonate, talc, titanium dioxide

Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg
 Cough suppressant
 Guaifenesin 200 mg
 Expectorant
 Phenylephrine HCl 5 mg
 Nasal decongestant

Active ingredients Purposes
(in each caplet)
Mucinex FAST-MAX
NIGHT TIME COLD & FLU
 Acetaminophen 325 mg

Pain reliever/fever reducer
 Diphenhydramine HCl 12.5 mg
 Antihistamine/cough suppressant
 Phenylephrine HCl 5 mg
 Nasal decongestant

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - sinus congestion and pressure
 - runny nose (**NIGHT TIME only**)
 - sneezing (**NIGHT TIME only**)
 - itching of the nose or throat (**NIGHT TIME only**)
 - itchy, watery eyes due to hay fever (**NIGHT TIME only**)
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**DAY TIME only**)
- controls cough to help you get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

Drug Facts (continued)

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin (**NIGHT TIME only**)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT TIME only**)
- a breathing problem such as emphysema or chronic bronchitis (**NIGHT TIME only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**NIGHT TIME only**)

When using this product

- **do not use more than directed**
- excitability may occur, especially in children (**NIGHT TIME only**)
- marked drowsiness may occur (**NIGHT TIME only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT TIME only**)
- avoid alcoholic drinks (**NIGHT TIME only**)
- be careful when driving a motor vehicle or operating machinery (**NIGHT TIME only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur



Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient


Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.


NDC 63824-548-64



Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant







Pediatrician Recommended!

TRIPLE PACK

READ ALL INFORMATION

Mucinex Children's

DAY TIME MULTI-SYMPTOM COLD

Dextromethorphan HBr 5 mg - Cough Suppressant
Guaifenesin 100 mg - Expectorant
Phenylephrine HCl 2.5 mg - Nasal Decongestant

✓ Relieves Stuffy Nose
✓ Controls Cough
✓ Relieves Chest Congestion
✓ Breaks up Mucus

DAY TIME MULTI-SYMPTOM COLD

Dextromethorphan HBr 5 mg - Cough Suppressant
Guaifenesin 100 mg - Expectorant
Phenylephrine HCl 2.5 mg - Nasal Decongestant

✓ Relieves Stuffy Nose
✓ Controls Cough
✓ Relieves Chest Congestion
✓ Breaks up Mucus

NIGHT TIME MULTI-SYMPTOM COLD

Acetaminophen 325 mg - Pain Reliever/Fever Reducer
Diphenhydramine HCl 12.5 mg - Antihistamine/Cough Suppressant
Phenylephrine HCl 5 mg - Nasal Decongestant

✓ Relieves Stuffy Nose
✓ Controls Cough
✓ Relieves Runny Nose & Sneezing
✓ Relieves Fever & Sore Throat

Very Berry Flavor Liquid

THREE - 4 FL OZ (118mL)

TOTAL - 12 FL OZ (354mL)

Print safe area

KO

Night Time Multi-Symptom Cold - Ages 6+

Drug Facts

Active Ingredients (in each 10 mL)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg.....Antihistamine/cough suppressant
Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

temporarily relieves these common cold and flu symptoms

headache
sore throat
nasal congestion
runny nose
sneezing
cough
controls cough to help your child get to sleep
temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blistering
- itching
- swelling
- fever
- hives
- difficulty breathing

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other product containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if the child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

Night Time Multi-Symptom Cold - Ages 6+ (cont'd)

Drug Facts (continued)**When using this product**

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or last more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

This product does not contain directions or complete warnings for adult use

- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20°-25°C (68°-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum, triethyl citrate, citric acid

Day Time Multi-Symptom Cold - Ages 4+

Drug Facts

Active ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg.....Cough suppressant
Guaifenesin 100 mg.....Expectorant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to clear the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- nasal congestion due to a cold
- stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get

Day Time Multi-Symptom Cold - Ages 4+ (cont'd)

Drug Facts (continued)

Other information

each 5 mL contains: sodium 3 mg
store at 20°-25°C (68°-77°F)
do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucrose, triethyl citrate, citric acid, xanthan gum
* may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

As an OTC brand in the
Children's Cough/Cold category
(AlphaImpactRxPreVoice Survey)

DO NOT GIVE THE MUCINEX CHILDREN'S DAY TIME MULTI-SYMPTOM COLD AND MUCINEX CHILDREN'S NIGHT TIME MULTI-SYMPTOM COLD LIQUIDS AT THE SAME TIME.

ALWAYS WAIT AT LEAST 4 HOURS BEFORE GIVING ANOTHER DOSE OF MUCINEX CHILDREN'S LIQUID. DO NOT GIVE MORE THAN A TOTAL OF 5 DOSES IN ANY 24-HOUR PERIOD.

Please visit our website www.mucinex.com

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Pediatrician Recommended!

TRIPLE PACK

Mucinex Children's

2X
DAY TIME
MULTI-SYMPTOM COLD

1X
NIGHT TIME
MULTI-SYMPTOM COLD

COUGH+ COLD NIGHT TIME – AGES 6+ (cont'd)

Drug Facts (continued)

When using this product:

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if:

- irritation, diarrhea, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup
- provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, fluoro, glycerin, propyl gallate, propylene glycol, purified water, sodium fumarate, sorbitol, sucralose, titanium dioxide, xanthan gum

may contain this ingredient

Keep cartons for full information.

COUGH+ CHEST CONGESTION DM – AGES 4+

Drug Facts

Active ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg..... Cough suppressant

Guaifenesin 100 mg..... Expectorant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has:

- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed.

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help and contact a Poison Control Center right away.

Directions

this product does not contain directions or complete warnings for adult use

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age (year)	Dose (mL)
children 6 years to under 12 years of age	5 mL – 10 mL every 4 hours
children 4 years to under 6 years of age	2.5 mL – 5 mL every 4 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

COUGH+ CHEST CONGESTION DM – AGES 4+ (cont'd)

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, fluoro, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, titanium dioxide, xanthan gum

may contain this ingredient

Questions? 1-800-862-4638

You may also report side effects to this phone number.

Tamper evident: Do not use if neckband on bottle cap is broken or missing

Do not give the Children's Delsym Cough+ Chest Congestion DM and Children's Delsym Cough+ Cold Night Time liquids at the same time.

Always wait at least 4 hours before giving another dose of Children's Delsym Cough+ Cold Night Time liquid. Do not give more than a total of 5 doses in any 24-hour period.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Please visit our website

www.delsym.com

Dist. by: RB Health (US)

Parity, NJ 07054-0224

© 2022 RB Health

Made in: US Patents: www.rb.com/patents

Day

children's Delsym

COMBO PACK

AGES 4+

COUGH+ CHEST CONGESTION DM

Dextromethorphan HBr (Cough Suppressant)

Guaifenesin (Expectorant)

- Relieves Cough & Chest Congestion
- Thins and Loosens Mucus

cherry

Flavored Liquid

Night

children's Delsym

COMBO PACK

AGES 6+

COUGH+ COLD NIGHT TIME

Acetaminophen (Pain Reliever/Fever Reducer)

Diphenhydramine HCl (Antihistamine/Cough Suppressant)

Phenylephrine HCl (Nasal Decongestant)

- Relieves Cough, Stuffy Nose & Sore Throat
- Reduces Fever

berry

Flavored Liquid

TWO - 4 fl oz (118 mL) bottles TOTAL – 8 fl oz (236 mL)

Drug Facts

Active ingredients (in each 10 mL)

Acetaminophen 325 mg...Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg...Antihistamine/cough suppressant
Phenylephrine HCl 5 mg...Nasal decongestant

Purposes

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
- temporarily reduces fever
- controls cough to help your child get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blistering
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other product containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if the child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

Drug Facts (continued)

When using this product

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or last more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient

Questions? 1-866-682-4639
You may also report side effects to this phone number.

Keep carton for full information.

**COUGH+
COLD
NIGHTTIME**

NDC 63824-212-64

**children's
Delsym®**

**AGES
6+**

**COUGH+
COLD
NIGHTTIME**

Acetaminophen (Pain Reliever/Fever Reducer)
Diphenhydramine HCl (Antihistamine/Cough Suppressant)
Phenylephrine HCl (Nasal Decongestant)

- Relieves Cough, Stuffy Nose & Sore Throat
- Reduces Fever

**berry
Flavored Liquid**

4 fl oz (118 mL)

reckitt

Dist. by: RB Health (US), Parsippany, NJ 07054-0224
©2022 RB Health 062322 3234388

Please visit our website
www.delsym.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Dosing cup included

1

1.00

Tamper evident: do not use if neckband on bottle cap is broken or missing.

Children's MucinEX®

Cold, Cough & Sore Throat

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Pain Reliever/Fever Reducer
Cough Suppressant
Expectorant
Nasal Decongestant

NDC 63824-278-64

Children's MucinEX®

Cold, Cough & Sore Throat

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Pain Reliever/Fever Reducer
Cough Suppressant
Expectorant
Nasal Decongestant

- ✓ Cough
- ✓ Sore Throat
- ✓ Nasal Congestion
- ✓ Chest Congestion

Age **6+**

Mixed Berry

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- **shake well before using**
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- children 6 to under 12 years of age: 10 mL in dosing cup provided
- children under 6 years of age: do not use

Other information

- each 10 mL contains: **sodium 6 mg**
- tamper evident: do not use if neckband on bottle cap is broken or missing.
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate

dihydrate, xanthan gum may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



4 FL OZ
(118 mL)



LOT CODE:

EXP. DATE:

3000454

Children's
Mucinex®

Drug Facts

Active ingredients
(in each 10 mL)

Purposes

Acetaminophen 325 mg...	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

■ temporarily relieves these common cold and

Cold, Cough & Sore Throat



DOSING CUP INCLUDED

Each dose of mixed berry-flavored Children's Mucinex® Cold, Cough & Sore Throat Liquid:

- ✓ Controls Cough
- ✓ Soothes Sore Throat
- ✓ Relieves Stuffy Nose
- ✓ Relieves Chest Congestion

Questions or comments?
1-866-MUCINEX (1-866-682-4639)
or www.mucinex.com

Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224 ©2013 RB

Reckitt Benckiser

Made in England 073013 3000454

- 2896
- It also relieves these common cold and flu symptoms:
- nasal congestion
 - stuffy nose
 - cough due to minor throat and bronchial irritation
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - minor aches and pains
 - sore throat
 - headache
 - temporarily reduces fever
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not use more than directed (see **Overdose warning**)

Tamper evident: do not use if neckband
on bottle cap is broken or missing.

Children's
MucinEX

Congestion & Cough

Guaifenesin 100 mg
Dextromethorphan HBr 5 mg
Phenylephrine HCl 2.5 mg

Expectorant
Cough Suppressant
Nasal Decongestant

VALUE SIZE!

NDC 63824-279-67

Children's
MucinEX

Congestion & Cough

Guaifenesin 100 mg
Dextromethorphan HBr 5 mg
Phenylephrine HCl 2.5 mg

Expectorant
Cough Suppressant
Nasal Decongestant

- ✓ Chest Congestion
- ✓ Breaks up Mucus
- ✓ Stuffy Nose
- ✓ Cough



**Berrylicious
Flavor Liquid**

Children's
MucinEX

Congestion & Cough



DOSING CUP INCLUDED

Each dose of berrylicious-flavored
**MucinEX® Congestion &
Cough Liquid:**

- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus
- ✓ Relieves Stuffy Nose



Controls Cough

Questions or comments?

1-866-MUCINEX (1-866-682-4639)
or **www.mucinex.com**



LOT CODE:

EXP. DATE:

8021397

Drug Facts (continued)

Inactive Ingredients anhydrous citric acid, D&C red #33, dextrose, FD&C blue #1, FD&C red #40, flavors, glycerin, methylparaben, potassium sorbate, propyl gallate, propylene glycol, propylparaben, purified water, saccharin sodium, sodium hydroxide, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Also look for

Drug Facts

Active Ingredients (In each 5 mL)	Purpose
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

Children's
MucinEX



**Great-tasting
chest-congestion relief in
fun-to-take, convenient
granule packets!**

Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224 ©2012 RB
Made in England 050112

 **Reckitt
Benckiser**
8021397

- heart disease
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma
- high blood pressure
- diabetes

When using this product

- do not use more than directed

Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other Information

- each 5 mL contains: sodium 3 mg
- tamper evident: do not use if neckband on bottle cap is broken or missing
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Tamper evident: do not use if neckband
on bottle cap is broken or missing.

Children's MucinEX®

Multi-Symptom Cold

Cough Suppressant
Expectorant
Nasal Decongestant

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg
Phenylephrine HCl 2.5 mg

VALUE SIZE!

NDC 63824-252-67

Children's MucinEX®

Multi-Symptom Cold

Cough Suppressant
Expectorant
Nasal Decongestant

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg
Phenylephrine HCl 2.5 mg

- ✓ Stuffy Nose
- ✓ Cough
- ✓ Chest Congestion
- ✓ Breaks up Mucus

Age **4+**

**Very Berry
Flavor Liquid**

6.8 FL OZ
(201 mL)

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Also look for
**Children's
MucinEX®**

mini-melts


Great-tasting
chest-congestion relief in
fun-to-take, convenient
granule packets!



Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224

Made in England ©2013 RB 072513 3000460







**Tamper evident: do not use if neckband
on bottle cap is broken or missing.**

**Children's
MucinEX**

**Multi-Symptom
Cold & Fever**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Pain Reliever/Fever Reducer

Cough Suppressant

Expectorant

Nasal Decongestant

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- **shake well before using**
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- children 6 to under 12 years of age: 10 mL in dosing cup provided
- children under 6 years of age: do not use

Other information

- each 10 mL contains: **sodium 6 mg**
- tamper evident: do not use if neckband on bottle cap is broken or missing.
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum *may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

NDC 63824-017-64

**Children's
MucinEX**

**Multi-Symptom
Cold & Fever**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Pain Reliever/Fever Reducer

Cough Suppressant

Expectorant

Nasal Decongestant

- ✓ **Fever**
- ✓ **Cough**
- ✓ **Stuffy Nose**
- ✓ **Chest Congestion**
- ✓ **Breaks up Mucus**

Age 6+



**Berry Blast
Flavor Liquid**

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

4 FL OZ
(118 mL)



3 63824 01764 8

LOT CODE:

EXP. DATE:

3000452

Children's
Mucinex

**Multi-Symptom
Cold & Fever**

Drug Facts

Active ingredients (in each 10 mL)

Purposes

Acetaminophen 325 mgPain reliever/fever reducer
Dextromethorphan HBr 10 mgCough suppressant
Guaifenesin 200 mgExpectorant
Phenylephrine HCl 5 mgNasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion ■ stuffy nose
 - cough due to minor throat and bronchial irritation
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - minor aches and pains
 - sore throat ■ headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial

COLD & FEVER



DOSING CUP INCLUDED

**Each dose of berry blast-flavored
Children's Mucinex®
Multi-Symptom
Cold & Fever Liquid:**

- ✓ **Reduces Fever**
- ✓ **Controls Cough**
- ✓ **Relieves Stuffy Nose**
- ✓ **Relieves Chest Congestion**
- ✓ **Breaks up Mucus**

Questions or comments?

**1-866-MUCINEX (1-866-682-4639)
or www.mucinex.com**

Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224 ©2013 RB



**Reckitt
Benckiser**

Made in England 072613 3000452

secretions to rid the bronchial passageways of
bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not use more than directed (see Overdose warning)

Tamper evident: do not use if neckband on bottle cap is broken or missing.

Children's
Mucinex
NIGHT TIME

Multi-Symptom Cold

New!

NDC 63824-600-64

Children's
Mucinex
NIGHT TIME

Multi-Symptom Cold
Pain Reliever/Fever Reducer
Antihistamine/Cough Suppressant
Nasal Decongestant

Acetaminophen 325 mg
Diphenhydramine HCl 12.5 mg
Phenylephrine HCl 5 mg

- ✓ Stuffy Nose
- ✓ Cough
- ✓ Runny Nose & Sneezing
- ✓ Fever & Sore Throat

Age **6+**

Mixed Berry
Flavor Liquid

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions


- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F)
- dosing cup provided ■ do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum *may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)



Multi-Symptom Cold



DOSING CUP INCLUDED

Each dose of mixed berry-flavored
**Children's Mucinex® Night Time
Multi-Symptom Cold Liquid:**

- ✓ Relieves Stuffy Nose
- ✓ Controls Cough
- ✓ Relieves Runny Nose & Sneezing
- ✓ Relieves Fever & Sore Throat

Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other drug containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed (see **Overdose warning**)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness ►

Tamper evident: do not use if neckband on bottle cap is broken or missing.

Children's Mucinex

Stuffy Nose & Cold

Expectorant
Nasal Decongestant
Guaifenesin 100 mg
Phenylephrine HCl 2.5 mg

Unvarnished & Unprinted Area

Drug Facts (continued)

Inactive Ingredients anhydrous citric acid, edetate disodium, FD&C red #40, flavors, glycerin, propylene glycol, propyl galate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum
*may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

NEW FORMULA

NDC 63824-284-64

Children's Mucinex

Stuffy Nose & Cold

Expectorant
Nasal Decongestant

Guaifenesin 100 mg
Phenylephrine HCl 2.5 mg

- ✓ Stuffy Nose
- ✓ Chest Congestion
- ✓ Breaks up Mucus

Children's Mucinex

Stuffy Nose & Cold

DOSING CUP INCLUDED

Each dose of mixed berry-flavored
Children's Mucinex®
Stuffy Nose & Cold Liquid:

- ✓ Relieves Stuffy Nose
- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus
- ✓ Alcohol-free

Questions or comments?
1-866-MUCINEX (1-866-682-4639)
or www.mucinex.com

Drug Facts

Active Ingredients (in each 5 mL) **Purpose**
Guaifenesin 100 mg..... Expectorant
Phenylephrine HCl 2.5 mg..... Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product

- do not use more than directed

Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache.

These could be signs of a serious illness.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- each 5 mL contains: sodium 3 mg
- tamper evident: do not use if neckband on bottle cap is broken or missing
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Also look for
Children's Mucinex

mini-melts

Great-tasting
chest-congestion relief in
fun-to-take, convenient
granule packets!

Reckitt Benckiser

Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in England ©2012 RB 060812 8034044

Mixed Berry
Flavor Liquid

4 FL OZ
(118 mL)

LOT CODE:
EXP. DATE:

8034044

Unvarnished & Unprinted Area

VALUE PACK
24 LIQUID GELS

MAXIMUM STRENGTH

Mucinex[®]
FAST-MAX[®]

DAY COLD & FLU **NIGHT COLD & FLU**

Tamper evident:
Do not use if carton is damaged or if printed seal on blister is broken or missing.

MAXIMUM STRENGTH

Mucinex[®]
FAST-MAX[®]

DAY COLD & FLU **NIGHT COLD & FLU**

***DAY COLD & FLU** helps to relieve these symptoms day or night; **NIGHT COLD & FLU** helps to relieve these symptoms at night

Maximum Strength per 4-hour dose

Do not take the Mucinex Fast-Max DAY COLD & FLU and Mucinex Fast-Max NIGHT COLD & FLU liquid gels at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid gels. Do not take more than a total of 12 liquid gels in any 24-hour period.

Take only as directed.

Keep carton for full information.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Fast Dissolving Liquid Gels!

MAXIMUM STRENGTH NDC 63824-519-24

Mucinex[®]
FAST-MAX[®]

DAY COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guafenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

NIGHT COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Doxylamine succinate – Antihistamine
Phenylephrine HCl – Nasal Decongestant


ALL IN ONE*

HEADACHE, SORE THROAT, FEVER, CHEST CONGESTION, COUGH, NASAL CONGESTION, SINUS CONGESTION, SINUS PRESSURE, BODY PAIN

ALL IN ONE*

HEADACHE, SORE THROAT, FEVER, ITCHY THROAT, COUGH, NASAL CONGESTION, RUNNY NOSE, SNEEZING, BODY PAIN

16 LIQUID GELS (Liquid Filled) **8 LIQUID GELS (Liquid Filled)** **24 TOTAL**



Drug Facts

Active ingredients (in each liquid gel) Purposes **Mucinex FAST-MAX DAY Cold & Flu**

Acetaminophen 325 mg...Pain reliever/fever reducer
Dextromethorphan HBr 10 mg...Cough suppressant
Guaifenesin 200 mg...Expectorant
Phenylephrine HCl 5 mg...Nasal decongestant

Active ingredients (in each liquid gel) Purposes **Mucinex FAST-MAX NIGHT Cold & Flu**

Acetaminophen 325 mg...Pain reliever/fever reducer
Dextromethorphan HBr 10 mg...Cough suppressant
Doxylamine succinate 6.25 mg...Antihistamine
Phenylephrine HCl 5 mg...Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:
■ sinus congestion and pressure
■ cough ■ minor aches and pains
■ headache ■ nasal congestion ■ sore throat
■ runny nose (NIGHT only) ■ sneezing (NIGHT only)
■ itching of the nose or throat (NIGHT only)
■ itchy, watery eyes due to hay fever (NIGHT only)
■ helps loosen phlegm (mucus) and thin bronchial secretions to
rid the bronchial passageways of bothersome mucus and
make coughs more productive (DAY only)
■ controls cough to help you get to sleep
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- glaucoma (NIGHT only)
- a breathing problem such as emphysema or chronic bronchitis (NIGHT only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers (NIGHT only)

When using this product ■ do not use more than directed
■ excitability may occur, especially in children (NIGHT only)
■ marked drowsiness may occur (NIGHT only)
■ alcohol, sedatives, and tranquilizers may increase drowsiness (NIGHT only)

■ avoid alcoholic drinks (NIGHT only)

■ be careful when driving a motor vehicle or operating machinery (NIGHT only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away.

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information ■ store at 20-25°C (68-77°F)
■ avoid excessive heat

Inactive ingredients (DAY only) FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Inactive ingredients (NIGHT only) D&C yellow no. 10,

Questions?

1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

www.mucinex.com

Patents: www.rb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
Made in China
©2019 RB Health
020519 8357061

MAXIMUM STRENGTH

Mucinex
FAST-MAX

DAY TIME NIGHT TIME
COLD & FLU

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

NDC 63824-511-40

MAXIMUM STRENGTH

Mucinex
FAST-MAX

DAY TIME
COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

NIGHT TIME
COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Diphenhydramine HCl – Antihistamine/
Cough Suppressant
Phenylephrine HCl – Nasal Decongestant



Actual Size

FOR AGES 12+

24 CAPLETS

TOTAL 40



Actual Size

FOR AGES 12+

16 CAPLETS

LOT:

EXP:

MADE IN:

3157569



Questions?
1-800-MUCINEX (1-800-882-4639)
You may also report side effects to this phone number

HEALTH • HYGIENE • HOME

www.mucinex.com
Patents: www.fda.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2020 RB Health

060220 3157569



3 63824 99511 3


MAXIMUM STRENGTH

Mucinex
FAST-MAX

DAY TIME NIGHT TIME
COLD & FLU

*DAY TIME COLD & FLU helps to relieve these symptoms: day or night.
NIGHT TIME COLD & FLU helps to relieve these symptoms at night.
Maximum Strength per 4-hour dose.
Do not take more than a total of 12 caplets in any 24-hour period.
Take only as directed.
Keep carton for full information.

PARENTS:
Learn about your medicine online
www.StopMedicineBlame.org



3138451

Do not take the Mucinex Fast-Max DAY TIME COLD & FLU and Mucinex Fast-Max NIGHT TIME COLD & FLU caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. Do not take more than a total of 12 caplets in any 24-hour period. Take only as directed.

Drug Facts

Active ingredients
(in each caplet)

Purposes

033120

Mucinex FAST-MAX DAY TIME COLD & FLU

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Active ingredients**Purposes****(in each caplet)****Mucinex FAST-MAX NIGHT TIME COLD & FLU**

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses**Mucinex Fast-Max DAY TIME COLD & FLU**

- temporarily relieves these common cold and flu symptoms:
 - cough ■ minor aches and pains ■ headache ■ nasal congestion
 - sore throat ■ sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- controls cough to help you get to sleep
- temporarily reduces fever

Mucinex Fast-Max NIGHT TIME COLD & FLU

- temporarily relieves these common cold and flu symptoms:
 - cough ■ minor aches and pains ■ headache ■ nasal congestion
 - sore throat ■ sinus congestion and pressure ■ runny nose ■ sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- controls cough to help you get to sleep
- temporarily reduces fever

Warnings**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. ▼

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Reseal Area**Drug Facts (continued)****Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin **(NIGHT TIME**

only)

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (*NIGHT TIME only*)
- a breathing problem such as emphysema or chronic bronchitis (*NIGHT TIME only*)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*NIGHT TIME only*)

When using this product

- do not use more than directed
- excitability may occur, especially in children (*NIGHT TIME only*)
- marked drowsiness may occur (*NIGHT TIME only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*NIGHT TIME only*)
- avoid alcoholic drinks (*NIGHT TIME only*)
- be careful when driving a motor vehicle or operating machinery (*NIGHT TIME only*)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. ▼

Hinge

Hinge

Drug Facts (continued)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

(Mucinex FAST-MAX DAY TIME COLD & FLU)

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients

(Mucinex FAST-MAX NIGHT TIME COLD & FLU)

croscarmellose sodium, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, ferric oxide yellow, methacrylic acid—ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium bicarbonate, talc, titanium dioxide

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



HEALTH ▸ HYGIENE ▸ HOME



Drug Facts (continued)**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

PEEL HERE

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 9 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

**Please visit our website
www.mucinex.com**

**Cooling
Menthol
Flavor!**

051017

3048117



For Multi-Symptom relief for both Day and Night, try our **Maximum Strength Mucinex® FAST-MAX® Day/Night packs**



HEALTH • HYGIENE • HOME

www.mucinex.com
Patents: www.rb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

Made in China
©2018 RB Health
020619 8357062

Drug Facts

Active ingredients (in each liquid gel)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - minor aches and pains
 - sinus congestion and pressure
 - temporarily reduces fever
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- nasal congestion
- cough
- sore throat

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure

Drug Facts (continued)

- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **Do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information


- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

PEEL HERE

more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided

Drug Facts (continued)

- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions?
1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.


HEALTH • HYGIENE • HOME

Please visit our website
www.mucinex.com







Drug Facts

Active ingredients (in each caplet) Purposes

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - cough
 - minor aches and pains
 - sore throat
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 12 caplets in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period



Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ liver disease ■ heart disease ■ diabetes

■ high blood pressure ■ thyroid disease

■ trouble urinating due to an enlarged prostate gland

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

Directions: Take 12 caplets in any 24-hour period.

■ adults and children 12 years of age and over: take

2 caplets every 4 hours

■ children under 12 years of age: do not use

Other information

■ store at 20-25°C (68-77°F)

Inactive ingredients croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Questions? 1-866-MUC INEX (1-866-682-4639)

You may also report side effects to this phone number.

Maximum Strength per 4h our dose
 *helps to relieve these symptoms day or night
 Do not take more than a total of 12 caplets in any 24-hour period.
 Take only as directed.
 Keep out of reach of children.

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org



Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over:
20 mL in dosing cup provided every 4 hours

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

MAXIMUM STRENGTH

Mucinex
FAST-MAX



NIGHT TIME COLD & FLU

NDC 63824-500-64

MAXIMUM STRENGTH

Mucinex
FAST-MAX



NIGHT TIME COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Diphenhydramine HCl – Antihistamine/Cough Suppressant
Phenylephrine HCl – Nasal Decongestant



■ children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate†, xanthan gum †may contain this ingredient

4 FL OZ (118 mL)

FOR AGES 12+



3 63824 19052 5

LOT CODE:

EXP. DATE:

MADE IN:

3100222

Pharma
Code

Keep carton for full information.

Maximum Strength per 4-hour dose

***Helps to relieve these symptoms at night**

Drug Facts

Active ingredients
(in each 20 mL)

Purposes

Acetaminophen 650 mg	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/ cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:

Questions? 1-866-Mucinex (1-866-682-4639)

You may also report side effects to this phone number.

Please visit our website www.mucinex.com

Patents: www.rb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2019 RB Health
042519 3100222



HEALTH • HYGIENE • HOME

- cough
- minor aches and pains
- headache
- sinus congestion and pressure
- runny nose
- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep
- nasal congestion
- sore throat
- sneezing

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- high blood pressure
- glaucoma